



Eidgenössische Technische Hochschule Zürich
Swiss Federal Institute of Technology Zurich

Forthcoming in: The Stata Journal

ETH Zurich Sociology Working Paper No. 5

A Stata implementation of the Blinder-Oaxaca decomposition

Ben Jann

May 2008

ETH Zurich, Chair of Sociology

SEW E 21, Scheuchzerstrasse 70
8092 Zurich, Switzerland

Tel. +41 44 632 55 56
Fax +41 44 632 10 54
info@soz.gess.ethz.ch
www.socio.ethz.ch

A Stata implementation of the Blinder-Oaxaca decomposition

Ben Jann

ETH Zurich
jannb@ethz.ch

May 2008

Abstract. The counterfactual decomposition technique popularized by Blinder (1973) and Oaxaca (1973) is widely used to study mean outcome differences between groups. For example, the technique is often used to analyze wage gaps by sex or race. The present paper summarizes the technique and addresses a number of complications such as the identification of effects of categorical predictors in the detailed decomposition or the estimation of standard errors. A new Stata command called `oaxaca` is introduced and examples illustrating its usage are given.

Keywords: st0001, Blinder-Oaxaca decomposition, outcome differential, wage gap

1 Introduction

An often used methodology to study labor market outcomes by groups (sex, race, and so on) is to decompose mean differences in log wages based on regression models in a counterfactual manner. The procedure is known in the literature as the Blinder-Oaxaca decomposition (Blinder 1973; Oaxaca 1973) and divides the wage differential between two groups into a part that is “explained” by group differences in productivity characteristics such as education or work experience and a residual part that cannot be accounted for by such differences in wage determinants. This “unexplained” part is often used as a measure for discrimination, but it also subsumes the effects of group differences in unobserved predictors. Most applications of the technique can be found in the labor market and discrimination literature (for meta studies see, e.g., Stanley and Jarrell 1998 or Weichselbaumer and Winter-Ebmer 2005). However, the method may also be useful in other fields. In general, the technique can be employed to study group differences in any (continuous and unbounded) outcome variable. For example, O’Donnell et al. (2008) use it to analyze health inequalities by poverty status.

The purpose of this article is to give an overview of the Blinder-Oaxaca decomposition and introduce a new command implementing the technique in Stata called `oaxaca`. In the next section, the most common variants of the procedure are summarized and a number of issues such as the identification of the contribution of categorical predictors or the estimation of standard errors are addressed. The third section then describes the syntax and options of the new `oaxaca` command and the fourth section provides examples for its application using labor market data.

2 Methods and formulas

Given are two groups A and B , an outcome variable Y , and a set of predictors. For example, think of males and females, (log) wages as the outcome variable, and human capital indicators such as education and work experience as predictors. The question now is how much of the mean outcome difference

$$R = E(Y_A) - E(Y_B) \quad (1)$$

where $E(Y)$ denotes the expected value of the outcome variable, is accounted for by group differences in the predictors.

Based on the linear model

$$Y_\ell = X'_\ell \beta_\ell + \epsilon_\ell, \quad E(\epsilon_\ell) = 0, \quad \ell \in \{A, B\} \quad (2)$$

where X is a vector containing the predictors and a constant, β contains the slope parameters and the intercept, and ϵ is the error, the mean outcome difference can be expressed as the difference in the linear prediction at the group-specific means of the regressors. That is

$$R = E(Y_A) - E(Y_B) = E(X_A)' \beta_A - E(X_B)' \beta_B \quad (3)$$

since

$$E(Y_\ell) = E(X'_\ell \beta_\ell + \epsilon_\ell) = E(X'_\ell \beta_\ell) + E(\epsilon_\ell) = E(X_\ell)' \beta_\ell$$

with $E(\beta_\ell) = \beta_\ell$ and $E(\epsilon_\ell) = 0$ by assumption.

To identify the contribution of group differences in predictors to the overall outcome difference, equation (3) can be rearranged, for example, as follows (see Winsborough and Dickinson 1971; Jones and Kelley 1984; Daymont and Andrisani 1984):

$$R = [E(X_A) - E(X_B)]' \beta_B + E(X_B)' (\beta_A - \beta_B) + [E(X_A) - E(X_B)]' (\beta_A - \beta_B) \quad (4)$$

This is a “three-fold” decomposition, that is, the outcome difference is divided into three parts:

$$R = E + C + I$$

The first summand

$$E = [E(X_A) - E(X_B)]' \beta_B$$

amounts to the part of the differential that is due to group differences in the predictors (the “endowments effect”). The second component

$$C = E(X_B)' (\beta_A - \beta_B)$$

measures the contribution of differences in the coefficients (including differences in the intercept). The third summand

$$I = [E(X_A) - E(X_B)]' (\beta_A - \beta_B)$$

is an interaction term accounting for the fact that differences in endowments and coefficients exist simultaneously between the two groups.

Decomposition (4) is formulated from the viewpoint of Group B . That is, the group differences in the predictors are weighted by the coefficients of Group B to determine the endowments effect (E). In other words, the E component measures the expected change in Group B 's mean outcome, if Group B had Group A 's predictor levels. Similarly, for the second component (C), the differences in coefficients are weighted by Group B 's predictor levels. That is, the second component measures the expected change in Group B 's mean outcome, if Group B had Group A 's coefficients. Naturally, the differential can analogously be expressed from the viewpoint of Group A , yielding the reverse three-fold decomposition

$$R = [E(X_A) - E(X_B)]' \beta_A + E(X_A)'(\beta_A - \beta_B) - [E(X_A) - E(X_B)]'(\beta_A - \beta_B) \quad (5)$$

Now the “endowments effect” amounts to the expected change of Group A 's mean outcome, if Group A had Group B 's predictor levels. The “coefficients effect” quantifies the expected change in Group A 's mean outcome, if Group A had Group B 's coefficients.

An alternative decomposition that is prominent in the discrimination literature results from the concept that there is some nondiscriminatory coefficients vector that should be used to determine the contribution of the differences in the predictors. Let β^* be such a nondiscriminatory coefficients vectors. The outcome difference can then be written as

$$R = [E(X_A) - E(X_B)]' \beta^* + [E(X_A)'(\beta_A - \beta^*) + E(X_B)'(\beta^* - \beta_B)] \quad (6)$$

We now have a “two-fold” decomposition

$$R = Q + U$$

where the first component

$$Q = [E(X_A) - E(X_B)]' \beta^*$$

is the part of the outcome differential that is “explained” by group differences in the predictors (the “quantity effect”) and the second summand

$$U = E(X_A)'(\beta_A - \beta^*) + E(X_B)'(\beta^* - \beta_B)$$

is the “unexplained” part. The latter is usually attributed to discrimination, but it is important recognize that it to also captures all potential effects of differences in unobserved variables.

The “unexplained” part in (6) is sometimes further decomposed. Let $\beta_A = \beta^* + \delta_A$ and $\beta_B = \beta^* + \delta_B$ with δ_A and δ_B as group-specific discrimination parameter vectors (positive or negative discrimination, depending on sign). U can then be expressed as

$$U = E(X_A)' \delta_A - E(X_B)' \delta_B \quad (7)$$

that is, the unexplained component of the differential can be subdivided into a part

$$U_A = E(X_A)' \delta_A$$

that measures discrimination in favor of Group *A* and a part

$$U_B = -E(X_B)' \delta_B$$

that quantifies discrimination against Group *B*.¹ Again, however, this interpretation hinges on the assumption that there are no relevant unobserved predictors.

The estimation of the components of the three-fold decompositions (4) and (5) is straightforward. Let $\hat{\beta}_A$ and $\hat{\beta}_B$ be the least squares estimates for β_A and β_B , obtained separately from the two group-specific samples. Furthermore, use the group means \bar{X}_A and \bar{X}_B as estimates for $E(X_A)$ and $E(X_B)$. Based on these estimates the decompositions (4) and (5) are computed as

$$\hat{R} = \bar{Y}_A - \bar{Y}_B = (\bar{X}_A - \bar{X}_B)' \hat{\beta}_B + \bar{X}_B' (\hat{\beta}_A - \hat{\beta}_B) + (\bar{X}_A - \bar{X}_B)' (\hat{\beta}_A - \hat{\beta}_B) \quad (8)$$

and

$$\hat{R} = \bar{Y}_A - \bar{Y}_B = (\bar{X}_A - \bar{X}_B)' \hat{\beta}_A + \bar{X}_A' (\hat{\beta}_A - \hat{\beta}_B) - (\bar{X}_A - \bar{X}_B)' (\hat{\beta}_A - \hat{\beta}_B) \quad (9)$$

The determination of the components of the two-fold decomposition (6) is more involved because an estimate for the unknown nondiscriminatory coefficients vector β^* is needed. Several suggestions have been made in the literature. For example, there may be reason to assume that discrimination is directed towards one of the groups only, so that $\beta^* = \beta_A$ or $\beta^* = \beta_B$ (see Oaxaca 1973, who speaks of an “index number problem”). Again assume that Group *A* are males and Group *B* are females. If, for instance, wage discrimination is only directed against women and there is no (positive) discrimination of men, then we may use $\hat{\beta}_A$ as an estimate for β^* and compute decomposition (6) as

$$\hat{R} = (\bar{X}_A - \bar{X}_B)' \hat{\beta}_A + \bar{X}_B' (\hat{\beta}_A - \hat{\beta}_B) \quad (10)$$

Similarly, if there is no discrimination of women, but only (positive) discrimination of men, the decomposition is

$$\hat{R} = (\bar{X}_A - \bar{X}_B)' \hat{\beta}_B + \bar{X}_A' (\hat{\beta}_A - \hat{\beta}_B) \quad (11)$$

Often, however, there is no specific reason to assume that the coefficients of one or the other group are nondiscriminating. Moreover, economists have argued that the undervaluation of one group comes along with an overvaluation of the other (e. g. Cotton 1988). Reimers (1983) therefore proposes to use the average coefficients over both groups as an estimate for the nondiscriminatory parameter vector, that is

$$\hat{\beta}^* = 0.5\hat{\beta}_A + 0.5\hat{\beta}_B \quad (12)$$

1. U_A and U_B have opposite interpretations. A positive value for U_A reflects *positive* discrimination of Group *A*; a positive value for U_B indicates *negative* discrimination of Group *B*.

Similarly, Cotton (1988) suggests to weight the coefficients by the group sizes n_A and n_B , that is

$$\hat{\beta}^* = \frac{n_A}{n_A + n_B} \hat{\beta}_A + \frac{n_B}{n_A + n_B} \hat{\beta}_B \quad (13)$$

Furthermore, based on theoretical derivations, Neumark (1988) advocates the usage of the coefficients from a pooled regression over both groups as an estimate for β^* .

Note that, as has been pointed out by Oaxaca and Ransom (1994) and others, decomposition (6) can also be expressed as

$$R = [E(X_A) - E(X_B)]' [\mathbf{W}\beta_A + (\mathbf{I} - \mathbf{W})\beta_B] + [(\mathbf{I} - \mathbf{W})'E(X_A) + \mathbf{W}'E(X_B)]' (\beta_A - \beta_B) \quad (14)$$

where \mathbf{W} is a matrix of relative weights given to the coefficients of Group A and \mathbf{I} is the identity matrix. For example, choosing $\mathbf{W} = \mathbf{I}$ is equivalent to setting $\beta^* = \beta_A$. Similarly, $\mathbf{W} = 0.5\mathbf{I}$ is equivalent to $\beta^* = 0.5\beta_A + 0.5\beta_B$. Furthermore, Oaxaca and Ransom (1994) show that

$$\widehat{\mathbf{W}} = \Omega = (\mathbf{X}'_A \mathbf{X}_A + \mathbf{X}'_B \mathbf{X}_B)^{-1} \mathbf{X}'_A \mathbf{X}_A \quad (15)$$

with \mathbf{X} as the observed data matrix, is equivalent to using the coefficients from a pooled model over both groups as the reference coefficients.²

An issue with the approach by Neumark (1988) or Oaxaca and Ransom (1994) is that it may inappropriately transfer some of the unexplained part of the differential into the explained component, although this does not seem to have received much attention in the literature.³ Assume a simple model of log wages ($\ln W$) on education (Z) with sex-specific intercepts α_M and α_F due to discrimination. The model is

$$\ln W = \begin{cases} \alpha_M + \gamma Z + \epsilon & \text{if "male"} \\ \alpha_F + \gamma Z + \epsilon & \text{if "female"} \end{cases} \quad (16)$$

Let $\alpha_M = \alpha$ and $\alpha_F = \alpha + \delta$, where δ is the discrimination parameter. Then the model may also be expressed as

$$\ln W = \alpha + \gamma Z + \delta F + \epsilon \quad (17)$$

with F as an indicator for "female". Assume that $\gamma > 0$ (positive relation between education and wages) and $\delta < 0$ (discrimination against women). If we now use γ^* from a pooled model

$$\ln W = \alpha^* + \gamma^* Z + \epsilon^* \quad (18)$$

in decomposition (6), then, following from the theory on omitted variables (see, e.g., Gujarati 2003, 510f.), the explained part of the differential is

$$Q = [E(Z_M) - E(Z_F)] \gamma^* = [E(Z_M) - E(Z_F)] \left(\gamma + \delta \frac{\text{Cov}(Z, G)}{\text{Var}(Z)} \right) \quad (19)$$

2. Another solution is to set $\mathbf{W} = \text{diag}(\beta - \beta_B) \cdot \text{diag}(\beta_A - \beta_B)^{-1}$, where β without subscript denotes the coefficients from the pooled model. Note that, although the decomposition results are the same, this approach yields a weighting matrix that is quite different from Oaxaca and Ransom's Ω . For example, whereas \mathbf{W} computed as described in this footnote is a diagonal matrix, Ω has off-diagonal elements unequal zero and is not even symmetric.

3. An exception is Fortin (2006)

with $Var(Z)$ as the variance of Z and $Cov(Z, G)$ as the covariance between Z and G . If men on average are better educated than women, then the covariance between Z and G is negative and the explained part of the decomposition gets overstated (given $\gamma > 0$ and $\delta < 0$). In essence, the difference in wages between men and women is explained by sex.

To avoid such a distortion of the decomposition results due to the residual group difference spilling over into the slope parameters of the pooled model, my recommendation is to always include a group indicator in the pooled model as an additional covariate.

Estimation of sampling variances

Given the popularity of the Blinder-Oaxaca procedure it is astonishing how little attention has been paid to the issue of statistical inference. Most studies in which the procedure is applied only report point estimates for the decomposition results, but do not make any indication about sampling variances or standard errors.⁴ However, for an adequate interpretation of the results approximate measures of statistical precision are indispensable.

Approximate variance estimators for certain variants of the decomposition were first proposed by Oaxaca and Ransom (1998) with Greene (2003, 53–54) making similar suggestions. The estimators by Oaxaca and Ransom (1998) and Greene (2003) are a good starting point, but they neglect an important source of variation. Most social-science studies on discrimination are based on survey data where all (or most) variables are random variables. That is, not only the outcome variable, but also the predictors are subject to sampling variation (an exception would be experimental factors set by the researcher). Whereas an important result for regression analysis is that it does not matter for the variance estimates whether regressors are stochastic or fixed, this is not true for the Blinder-Oaxaca decomposition. The decomposition is based on multiplying regression coefficients by means of regressors. If the regressors are stochastic, then the means have sampling variances. These variances are of the same asymptotic order than the variances of the coefficients (think of the means as the intercepts from regression models without covariates). To get consistent standard errors for the decomposition results it seems therefore important to take the variability induced by the randomness of the predictors into account.

Consider the expression

$$\bar{Y} = \bar{X}'\hat{\beta} \quad (20)$$

where \bar{X} is the vector of mean estimates for the predictors and $\hat{\beta}$ contains the least-squares estimates of the regression coefficients. If the predictors are stochastic, then \bar{X} and $\hat{\beta}$ are both subject to sampling variation. Assuming that \bar{X} and $\hat{\beta}$ are uncorrelated (which follows from the standard regression assumption that the conditional expectation of the error is zero for all covariate values; of course this is only true if the model is

4. Exceptions are for example Oaxaca and Ransom (1994, 1998), Silber and Weber (1999), Horrace and Oaxaca (2001), Fortin (2006), Heinrichs and Kennedy (2007) and Lin (2007). Furthermore, Jackson and Lindley (1989) and Shrestha and Sakellariou (1996) propose statistical tests for discrimination.

correctly specified), the variance of (20) can be written as

$$V(\bar{X}'\hat{\beta}) = E(\bar{X})'V(\hat{\beta})E(\bar{X}) + E(\hat{\beta})'V(\bar{X})E(\hat{\beta}) + \text{trace} \left[V(\bar{X})V(\hat{\beta}) \right] \quad (21)$$

where $V(\bar{X})$ and $V(\hat{\beta})$ are the variance-covariance matrices for \bar{X} and $\hat{\beta}$ (see the proof in Jann 2005b; for the variance of the product of two independent random variables also see Mood et al. 1974, 180). By inserting estimates for the expectations and variance matrices, we get the variance estimator

$$\widehat{V}(\bar{X}'\hat{\beta}) = \bar{X}'\widehat{V}(\hat{\beta})\bar{X} + \hat{\beta}'\widehat{V}(\bar{X})\hat{\beta} + \text{trace} \left[\widehat{V}(\bar{X})\widehat{V}(\hat{\beta}) \right] \quad (22)$$

$\widehat{V}(\hat{\beta})$ is simply the variance-covariance matrix obtained from the regression procedure. A natural estimator for $V(\bar{X})$ is $\widehat{V}(\bar{X}) = \mathcal{X}'\mathcal{X}/[n(n-1)]$, where \mathcal{X} is the centered data matrix, i. e. $\mathcal{X} = \mathbf{X} - \mathbf{1}\bar{X}'$.

The variances for the components of the Blinder-Oaxaca decomposition can be derived analogously. For example, ignoring the asymptotically vanishing⁵ last term in (22) and assuming that the two groups are independent, the approximate variance estimators for the two terms of decomposition (10) are

$$\widehat{V}([\bar{X}_A - \bar{X}_B]'\hat{\beta}_A) \approx (\bar{X}_A - \bar{X}_B)'\widehat{V}(\hat{\beta}_A)(\bar{X}_A - \bar{X}_B) + \hat{\beta}_A' \left[\widehat{V}(\bar{X}_A) + \widehat{V}(\bar{X}_B) \right] \hat{\beta}_A \quad (23)$$

and

$$\widehat{V}(\bar{X}_B'[\hat{\beta}_A - \hat{\beta}_B]) \approx \bar{X}_B' \left[\widehat{V}(\hat{\beta}_A) + \widehat{V}(\hat{\beta}_B) \right] \bar{X}_B + (\hat{\beta}_B - \hat{\beta}_B)'\widehat{V}(\bar{X}_B)(\hat{\beta}_B - \hat{\beta}_B) \quad (24)$$

where use of the fact is made that the variance of the sum of two uncorrelated random variables is equal to the sum of the individual variances. An interesting point about formulas (23) and (24) is that ignoring the stochastic nature of the predictors will primarily affect the variance of the first term of the decomposition (the “explained” part). This is because in most applications group differences in coefficients and means are much smaller than the levels of coefficients and means.

It is possible to develop similar formulas for all the decomposition variants outlined above, but derivations can get complicated once a pooled model is used and covariances between the pooled model and the group models have to be taken into account or if the assumption of independence between the two groups is loosened (e. g. if dealing with a cluster sample). An alternative approach that is simple and general and produces equivalent results is to estimate the joint variance-covariance matrix of all used statistics (see Weesie 1999, [R] **suest**) and then apply the “delta method” (see [R] **nlcom** and the references therein). In fact, in the case of independence between the two groups the results of the delta method for decomposition (4) are formally equal to (23) and (24). Furthermore, a general result for the delta method is that if the input variance matrix is asymptotically normal, then also the variance matrix of the transformed statistics is

5. Whereas the first and second terms are of order $O(n^{-1})$, the last term is $O(n^{-2})$.

asymptotically normal (see, e. g., Greene 2003, 70). That is, since asymptotic normality holds for regression coefficients and mean estimates under very general conditions, the variances obtained by the delta method can be used to construct approximate confidence intervals for the decomposition results in the usual manner.

Detailed decomposition

Often not only is the total decomposition of the outcome differential into an explained and an unexplained part of interest, but also the detailed contributions of the single predictors or sets of predictors are subject to investigation. For example, one might want to evaluate how much of the gender wage gap is due to sex differences in education and how much is due to differences in work experience. Similarly, it might be informative to determine how much of the unexplained gap is related to differing returns to education and how much to differing returns to work experience.

Identifying the contributions of the individual predictors to the explained part of the differential is easy because the total component is a simple sum over the individual contributions. For example, for decomposition (10),

$$\widehat{Q} = (\bar{X}_A - \bar{X}_B)' \hat{\beta}_A = (\bar{X}_{1A} - \bar{X}_{1B}) \hat{\beta}_{1A} + (\bar{X}_{2A} - \bar{X}_{2B}) \hat{\beta}_{2A} + \dots \quad (25)$$

where $\bar{X}_1, \bar{X}_2, \dots$ are the means of the single regressors and $\hat{\beta}_1, \hat{\beta}_2, \dots$ are the associated coefficients. The first summand reflects the contribution of the group differences in \bar{X}_1 , the second of differences in \bar{X}_2 , and so on. Also the estimation of standard errors for the individual contributions is straightforward.

Similarly, using decomposition (10) as an example, the individual contributions to the unexplained part are the summands in

$$\widehat{U} = \bar{X}'_B (\hat{\beta}_A - \hat{\beta}_B) = \bar{X}'_{1B} (\hat{\beta}_{1A} - \hat{\beta}_{1B}) + \bar{X}'_{2B} (\hat{\beta}_{2A} - \hat{\beta}_{2B}) + \dots \quad (26)$$

Note, however, that other than for the explained part of the decomposition, the contributions to the unexplained part may depend on arbitrary scaling decisions if the predictors do not have natural zero points (e. g. Jones and Kelley 1984, 334). Without loss of generality, assume a simple model with just one explanatory variable:

$$Y_\ell = \beta_{0\ell} + \beta_{1\ell} Z_\ell + \epsilon_\ell, \quad \ell \in \{A, B\}$$

The unexplained part of the decomposition based on (10) then is

$$\widehat{U} = (\hat{\beta}_{0A} - \hat{\beta}_{0B}) + (\hat{\beta}_{1A} - \hat{\beta}_{1B}) \bar{Z}_B$$

The first summand is the part of the unexplained gap that is due to “group membership” (Jones and Kelley 1984); the second summand reflects the contribution of differing returns to Z . Now assume that the zero point of Z is shifted by adding a constant a . The effect of such a shift on the decomposition results is as follows:

$$\widehat{U} = \left[(\hat{\beta}_{0A} - a\hat{\beta}_{1A}) - (\hat{\beta}_{0B} - a\hat{\beta}_{1B}) \right] + (\hat{\beta}_{1A} - \hat{\beta}_{1B})(\bar{Z}_B + a)$$

Evidently, the scale shift changes the results; a portion amounting to $a(\hat{\beta}_{1A} - \hat{\beta}_{1B})$ is transferred from the group membership component to the part that is due to different slope coefficients. The conclusion is that the detailed decomposition results for the unexplained part only have a meaningful interpretation for variables for which scale shifts are not allowed, that is, for variables that have a natural zero point.⁶

A related issue that has received much attention in the literature is that the decomposition results for categorical predictors depend on the choice of the omitted base category (Jones 1983; Jones and Kelley 1984; Oaxaca and Ransom 1999; Nielsen 2000; Horrace and Oaxaca 2001; Gardeazabal and Ugidos 2004; Polavieja 2005; Yun 2005b). The effect of a categorical variable is usually modeled by including 0/1-variables (“dummy” variables) for the different categories in the regression equation, where one of the categories (the “base” category) is omitted to avoid collinearity. It is easy to see that the decomposition results for the single 0/1-variables depend on the choice of the base category, since the associated coefficients quantify differences with respect to the base category. If the base category changes, the decomposition results change.

For the explained part of the decomposition this may not be critical because the sum of the contributions of the single indicator variables, that is, the total contribution of the categorical variable is unaffected by the choice of the base category. For the unexplained part of the decomposition, however, there is again a tradeoff between the group membership component (the difference in intercepts) and the part attributed to differences in slope coefficients. For the unexplained part changing the base category not only alters the results for the singly dummy variables but also changes the contribution of the categorical variable as a whole.

An intuitively appealing solution to the problem has been proposed by Gardeazabal and Ugidos (2004) and Yun (2005b). The idea is to restrict the coefficients for the single categories to sum to zero, that is, to express effects as deviations from the grand mean. This can be implemented by restricted least squares estimation or by transforming the dummy variables before model estimation as proposed by Gardeazabal and Ugidos (2004).⁷ A more convenient method in the context of the Blinder-Oaxaca decomposition is to estimate the group models using the standard dummy coding and then transform the coefficients vectors so that deviations from the grand mean are expressed and the (redundant) coefficient for the base category is added (Suits 1984; Yun 2005b). If applied to such transformed estimates, the results of the Blinder-Oaxaca decomposition are independent of the choice of the omitted category. Furthermore, the results are equal to the simple averages of the results one would get from a series of decompositions in which the categories are used one after another as the base category (Yun 2005b).

6. The problem does not occur for the explained part of the decomposition or the interaction component in the three-fold decomposition because a cancels out in these cases. Furthermore, stretching or compressing the scales of the X variables (multiplication by a constant) does not alter any of the decomposition results because such multiplicative transformations are counterbalanced by the coefficient estimates.

7. In fact, the approach by Gardeazabal and Ugidos (2004) is simply what is known as the “effects coding” (Hardy 1993, 64–71) or the “deviation contrast coding” (Hendrickx 1999).

The deviation contrast transform works as follows. Given is the model

$$Y = \beta_0 + \beta_1 D_1 + \dots + \beta_{k-1} D_{k-1} + \epsilon$$

where β_0 is the intercept and D_j , $j = 1, \dots, k-1$, are the dummy variables representing a categorical variable with k categories. Category k is the base category. Alternatively, the model may be formulated as

$$Y = \beta_0 + \beta_1 D_1 + \dots + \beta_{k-1} D_{k-1} + \beta_k D_k + \epsilon$$

where β_k is constrained to zero. Now let

$$c = (\beta_1 + \dots + \beta_k)/k$$

and define

$$\tilde{\beta}_0 = \beta_0 + c \quad \text{and} \quad \tilde{\beta}_j = \beta_j - c, \quad j = 1, \dots, k$$

The transformed model then is

$$Y = \tilde{\beta}_0 + \tilde{\beta}_1 D_1 + \dots + \tilde{\beta}_k D_k + \epsilon, \quad \sum_{j=1}^k \tilde{\beta}_j = 0$$

Note that the transformed model is mathematically equivalent to the untransformed model. For example, the two models produce identical predictions. The variance-covariance matrix for the transformed model can be obtained by applying the general formula for weighted sums of random variables given in, e.g., Mood et al. (1974, 179). Models with several sets of dummy variables can be transformed by applying the formulas to each set separately. Furthermore, the transformation can be applied to the interaction terms between a categorical and a continuous variable in an analogous manner except that now c is added to the main effect of the continuous variable instead of the intercept. Also note that the application of the transform is not restricted to linear regression. It can be used with any model as long as the effects of the dummies are expressed as additive effects.

Other restrictions to identify the contribution of a categorical variable to the unexplained part of the decomposition are imaginable. For example, the restriction could be

$$\sum_{j=1}^k w_j \tilde{\beta}_j = 0$$

where w_j are weights proportional to the relative frequencies of the categories, so that the coefficients reflect deviations from the overall sample mean (Kennedy 1986; Haisken-DeNew and Schmidt 1997). Hence, there is still some arbitrariness in the method by Gardeazabal and Ugidos (2004) and Yun (2005b).

3 The `oaxaca` command

The methods presented above are implemented in a new command called `oaxaca`. The command first estimates the group models and possibly a pooled model over both

groups using `regress` ([R] `regress`) or any user specified estimation command. `suest` is then applied, if necessary, to determine the combined variance-covariance matrix of the models ([R] `suest`) and the group means of the predictors are estimated using `mean` ([R] `mean`). Finally, the various decomposition results and their standard errors (and covariances) are computed based on the combined parameter vector and variance-covariance matrix of the models' coefficients and the mean estimates.⁸ The standard errors are obtained by the delta method.⁹

3.1 Syntax

The syntax of the `oaxaca` command is

```
oaxaca depvar [ indepvars ] [ if ] [ in ] [ weight ], by(groupvar) [ swap
  detail[(dlist)] adjust(varlist) threefold[(reverse)] weight(# [# ...])
  pooled[(model_opts)] omega[(model_opts)] reference(name) split
  x1(names_and_values) x2(names_and_values) categorical(clist)
  svy[(svyspec)] vce(vctype) cluster(varname) fixed[(varlist)] [no]suest
  nose model1(model_opts) model2(model_opts) xb noisily level(#) eform ]
```

where `depvar` is the outcome variable of interest (e.g. log wages) and `indepvars` are predictors (e.g. education, work experience, etc.). `groupvar` identifies the groups to be compared. `oaxaca` typed without arguments replays the last results.

`fweights`, `aweight`s, `pweight`s, and `iweight` are allowed; see [U] **11.1.6 weight**. Furthermore, `bootstrap`, `by`, `jackknife`, `statsby`, and `xi` are allowed; see [U] **11.1.10 Prefix commands**. Weights are not allowed with the `bootstrap` prefix and `aweight`s are not allowed with the `jackknife` prefix. `vce()`, `cluster()`, and weights are not allowed with the `svy` option.

3.2 Options

Main

`by(groupvar)` specifies the `groupvar` that defines the two groups that will be compared. `by()` is not optional.

`swap` reverses the order of the groups.

`detail[(dlist)]` requests that the detailed results for the individual predictors be re-

8. The covariances between the models' coefficients and the mean estimates are assumed zero in any case. This assumption may be violated in misspecified models.

9. `nlcom` ([R] `nlcom`) could be used to compute the variance-covariance matrix of the decomposition results. However, `nlcom` employs general methods based on numerical derivatives and is slow if the models contain many covariates. `oaxaca` therefore has its own specific implementation of the delta method based on analytic derivatives.

ported. Use *dlist* to subsume the results for sets of regressors (variables not appearing in *dlist* are listed individually). The syntax for *dlist* is

```
name:varlist [, name:varlist ...]
```

The usual shorthand conventions apply to the *varlists* specified in *dlist* (see help *varlist*; additionally, `_cons` is allowed). For example, specify `detail(exp:exp*)` to subsume `exp` (experience) and `exp2` (experience squared). *name* is any valid Stata name and labels the set.

`adjust(varlist)` causes the differential to be adjusted by the contribution of the specified variables before performing the decomposition. This is useful, for example, if the specified variables are selection terms. Note that `adjust()` is not needed for `heckman` models.

Decomposition type

`threefold[(reverse)]` computes the three-fold decomposition. This is the default unless `weight()`, `pooled`, `omega`, or `reference()` is specified. The decomposition is expressed from the viewpoint of Group 2 (*B*). Specify `threefold(reverse)` to express the decomposition from the viewpoint of Group 1 (*A*).

`weight(# [# ...])` computes the two-fold decomposition where `# [# ...]` are the weights given to Group 1 (*A*) relative to Group 2 (*B*) in determining the reference coefficients (weights are recycled if there are more coefficients than weights). For example, `weight(1)` uses the Group 1 coefficients as the reference coefficients, `weight(0)` uses the Group 2 coefficients.

`pooled[(model_opts)]` computes the two-fold decomposition using the coefficients from a pooled model over both groups as the reference coefficients. *groupvar* is included in the pooled model as an additional control variable. Estimation details may be specified in parentheses; see the `model11()` option below.

`omega[(model_opts)]` computes the two-fold decomposition using the coefficients from a pooled model over both groups as the reference coefficients (without including *groupvar* as a control variable in the pooled model). Estimation details may be specified in parentheses; see the `model11()` option below.

`reference(name)` computes the two-fold decomposition using the coefficients from a stored model. *name* is the name under which the model was stored; see [R] `estimates store`. Do not combine the `reference()` option with bootstrap or jackknife methods.

`split` causes the "unexplained" component in the two-fold decomposition to be split into a part related to Group 1 (*A*) and a part related to Group 2 (*B*). `split` is effective only if specified with `weight()`, `pooled`, `omega`, or `reference()`.

Only one of `threefold`, `weight()`, `pooled`, `omega`, and `reference()` is allowed.

X-Values

`x1(names_and_values)` and `x2(names_and_values)` provide custom values for specific predictors to be used for Group 1 (*A*) and Group 2 (*B*) in the decomposition. The default is to use the group means of the predictors. The syntax for *names_and_values* is

```
varname [=] value [[,] varname [=] value ... ]
```

Example: `x1(educ 12 exp 30)`

`categorical(clist)` identifies sets of dummy variables representing categorical variables and transforms the coefficients so that the results of the decomposition are invariant to the choice of the (omitted) base category (deviation contrast transform). The syntax for *clist* is

```
varlist [, varlist ... ]
```

Each variable list must contain a variable for the base category (that is, the base category indicator must exist in the data). The transform can also be applied to interactions between a categorical and a continuous variable. Specify the continuous variable in parentheses at the end of the list in this case, i.e.

```
varlist (varname) [, ... ]
```

and also include a list for the main effects. Example:

```
detail(d1 d2 d3, xd1 xd2 xd3 (x))
```

where *x* is the continuous variable, and *d1* etc. and *xd1* etc. are the main effects and interaction effects. The code for implementing the `categorical()` option has been taken from the `devcon` user command (Jann 2005a).

SE/SVY

`svy`[[*vcetype*] [, *svy_options*]] executes `oaxaca` while accounting for the survey settings identified by `svyset` (this is essentially equivalent to applying the `svy` prefix command, although the `svy` prefix is not allowed with `oaxaca` due to some technical issues). *vcetype* and *svy_options* are as described in [SVY] `svy`.

`vce`(*vcetype*) specifies the type of standard errors reported. *vcetype* may be `analytic` (the default), `robust`, `cluster clustvar`, `bootstrap`, or `jackknife`; see [R] *vce_option*.

`cluster`(*varname*) adjusts standard errors for intragroup correlation; this is Stata 9 syntax for `vce(cluster clustvar)`.

`fixed`[[*varlist*]] identifies fixed regressors (all if specified without argument; an example for fixed regressors are experimental factors). The default is to treat regressors as stochastic. Stochastic regressors inflate the standard errors of the decomposition components.

[no]suest prevents or enforces using suest to obtain the covariances between the models/groups. suest is implied by pooled, omega, reference(), svy, vce(cluster), and cluster(). Specifying nosuest may cause biased standard errors and is strongly discouraged.

nose suppresses the computation of standard errors.

Model estimation

model1(*model_opts*) and model2(*model_opts*) specify the estimation details for the two group-specific models. The syntax for *model_opts* is

```
[estcom] [, addrhs(spec) estcom_options ]
```

where *estcom* is the estimation command to be used and *estcom_options* are options allowed by *estcom*. The default estimation command is **regress**. **addrhs**(*spec*) adds *spec* to the “right-hand side” of the model. For example, use **addrhs**() to add extra variables to the model. Examples:

```
model1(heckman, select(varlist_s) twostep)
model1(ivregress 2sls, addrhs((varlist2=varlist_iv)))
```

Note that **oaxaca** uses the first equation for the decomposition if a model contains multiple equations.

Furthermore, coefficients that occur in one of the model only are assumed zero for the other group. It is important, however, that the associated variables contain non-missing values for all observations in both groups.

noisily displays the models’ estimation output.

Reporting

xb displays a table containing the regression coefficients and predictor values on which the decomposition is based.

level(#) specifies the confidence level, as a percentage, for confidence intervals. The default is **level**(95) or as set by **set level**.

eform specifies that the results be displayed in exponentiated form.

3.3 Returned results

Scalars

`e(N)` number of observations `e(N_clust)` number of clusters
`e(N1)` number of obs. in Group 1 `e(N1)` number of obs. in Group 2

Macros

`e(cmd)` `oaxaca`
`e(depvar)` name of dependent variable `e(by)` name group variable
`e(group1)` value defining Group 1 `e(group2)` value defining Group 2
`e(wtype)` weight type `e(wexp)` weight expression
`e(clustvar)` name of cluster variable `e(vce)` *vce*type specified in `vce()`
`e(vcetype)` title used to label Std. Err. `e(properties)` `b V`

Matrices

`e(b)` decomposition results `e(V)` variance matrix of `e(b)`
`e(b0)` coefficients and *X*-values `e(V0)` variance matrix of `e(b0)`

Functions

`e(sample)` marks estimation sample

4 Examples

Three-fold decomposition

The standard application of the Blinder-Oaxaca technique is to divide the wage gap between, say, men and women into a part that is explained by differences in determinants of wages such as education or work experience and a part that cannot be explained by such group differences. An example using data from the Swiss Labor Market Survey 1998 (Jann 2003) is as follows:

```
. use oaxaca, clear
(Excerpt from the Swiss Labor Market Survey 1998)
. oaxaca lnwage educ exper tenure, by(female) noisily
```

Model for group 1

Source	SS	df	MS			
Model	49.613308	3	16.5377693	Number of obs =	751	
Residual	122.143834	747	.163512495	F(3, 747) =	101.14	
				Prob > F =	0.0000	
				R-squared =	0.2889	
				Adj R-squared =	0.2860	
Total	171.757142	750	.229009522	Root MSE =	.40437	

lnwage	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
educ	.0820549	.0060851	13.48	0.000	.070109	.0940008
exper	.0098347	.0016665	5.90	0.000	.0065632	.0131062
tenure	.0100314	.0020397	4.92	0.000	.0060272	.0140356
_cons	2.24205	.0778703	28.79	0.000	2.08918	2.394921

Model for group 2

Source	SS	df	MS			
Model	33.5197344	3	11.1732448	Number of obs =	683	
Residual	188.08041	679	.276996185	F(3, 679) =	40.34	
				Prob > F =	0.0000	
				R-squared =	0.1513	
				Adj R-squared =	0.1475	
Total	221.600144	682	.324926897	Root MSE =	.5263	

be used, but now the pooled models also contains a group membership indicator. Based on the argumentation outlined in Section 2 my suggestion is to use `pooled` instead of `omega`.

For our example data, the results with the `pooled` option are as follows:

```
. oaxaca lnwage educ exper tenure, by(female) pooled
Blinder-Oaxaca decomposition          Number of obs   =       1434
      1: female = 0
      2: female = 1
```

	Coef.	Robust Std. Err.	z	P> z	[95% Conf. Interval]	
Differential						
Prediction_1	3.440222	.0174586	197.05	0.000	3.406004	3.47444
Prediction_2	3.266761	.0218042	149.82	0.000	3.224026	3.309497
Difference	.1734607	.0279325	6.21	0.000	.118714	.2282075
Decomposit-n						
Explained	.089347	.0137531	6.50	0.000	.0623915	.1163026
Unexplained	.0841137	.025333	3.32	0.001	.034462	.1337654

Again the conclusion is that differences in endowments account for about half of the wage gap.¹⁰

A further possibility is to provide a stored reference model using the `reference()` option. For example, for the decomposition of the wage gap between blacks and whites the reference model is sometimes estimated based on all races, not just blacks and whites. In such a case, the reference model would have to be estimated first using all observations and then be provided to `oaxaca` via the `reference()` option.

Exponentiated results

The results in the example above are expressed on the logarithmic scale (remember that log wages are used as the dependent variable) and it might be sensible to re-transform the results to the original scale (Swiss francs in this case) using the `eform` option:

```
. oaxaca, eform
Blinder-Oaxaca decomposition          Number of obs   =       1434
      1: female = 0
      2: female = 1
```

	exp(b)	Robust Std. Err.	z	P> z	[95% Conf. Interval]	
--	--------	---------------------	---	------	----------------------	--

10. Note that unlike in the first example robust standard errors are reported (`oaxaca` uses `suest` to estimate the joint variance matrix for all coefficients if `pooled` is specified; `suest` implies robust standard errors). To compute robust standard errors in the first example you would have to add `vce(robust)` to the command.

Differential						
Prediction_1	31.19388	.5446007	197.05	0.000	30.14454	32.27975
Prediction_2	26.22626	.5718438	149.82	0.000	25.12908	27.37135
Difference	1.189414	.0332234	6.21	0.000	1.126048	1.256346
Decomposit-n						
Explained	1.09346	.0150385	6.50	0.000	1.064379	1.123336
Unexplained	1.087753	.027556	3.32	0.001	1.035063	1.143125

The (geometric) means of wages are 31.2 Swiss francs for men and 26.2 francs for women, which amounts to a difference of 18.9 percent. Adjusting women's endowments levels to the levels of men would increase women's wages by 9.3 percent. A gap of 8.8 percent remains unexplained.

Survey estimation

`oaxaca` supports complex survey estimation, but `svy` has to be specified as an option and is not allowed as a prefix command (which does not restrict functionality). For example, variable `wt` provides sampling weights for the Swiss Labor Market Survey 1998. The weights (and strata or PSU's if there were any) can be taken into account as follows:

```
. svyset [pw=wt]
      pweight: wt
      VCE: linearized
      Single unit: missing
      Strata 1: <one>
      SU 1: <observations>
      FPC 1: <zero>

. oaxaca lnwage educ exper tenure, by(female) pooled svy
Blinder-Oaxaca decomposition
Number of strata =          1          Number of obs   =       1647
Number of PSUs   =       1434          Population size = 1424.3797
                                          Design df      =       1433

      1: female = 0
      2: female = 1
```

	Linearized		t	P> t	[95% Conf. Interval]	
	Coef.	Std. Err.				
Differential						
Prediction_1	3.405696	.0226315	150.48	0.000	3.361302	3.450091
Prediction_2	3.193847	.0276466	115.52	0.000	3.139615	3.248079
Difference	.2118488	.0357284	5.93	0.000	.1417633	.2819344
Decomposit-n						
Explained	.1107614	.0189974	5.83	0.000	.0734956	.1480271
Unexplained	.1010875	.0315911	3.20	0.001	.0391178	.1630572

Detailed decomposition

Use the `detail` option to compute the individual contributions of the predictors to the components of the decomposition. `detail` specified without argument reports the contribution of each predictor individually. Alternatively, one can define groups of predictors for which the results be subsumed in parentheses. Furthermore, one might apply the deviation contrast transform to dummy variable sets so that the contribution of a categorical predictor to the unexplained part of the decomposition does not depend on the choice of the base category. Example:

```
. tabulate isco, nofreq generate(isco)
. oaxaca lnwage educ exper tenure isco2-isco9, by(female) pooled ///
>   detail(exp_ten: exper tenure, isco: isco?) categorical(isco?)
Blinder-Oaxaca decomposition          Number of obs   =      1434
      1: female = 0
      2: female = 1
```

	Coef.	Robust Std. Err.	z	P> z	[95% Conf. Interval]	
Differential						
Prediction_1	3.440222	.0174589	197.05	0.000	3.406003	3.474441
Prediction_2	3.266761	.0218047	149.82	0.000	3.224025	3.309498
Difference	.1734607	.0279331	6.21	0.000	.118713	.2282085
Explained						
educ	.0395615	.0097334	4.06	0.000	.0204843	.0586387
exp_ten	.0399316	.0089081	4.48	0.000	.022472	.0573911
isco	-.0056093	.012445	-0.45	0.652	-.0300009	.0187824
Total	.0738838	.017772	4.16	0.000	.0390513	.1087163
Unexplained						
educ	-.1324971	.1788045	-0.74	0.459	-.4829475	.2179533
exp_ten	.0129955	.0400811	0.32	0.746	-.0655619	.0915529
isco	-.0159367	.0296549	-0.54	0.591	-.0740592	.0421858
_cons	.2350152	.195018	1.21	0.228	-.1472132	.6172435
Total	.0995769	.0266887	3.73	0.000	.047268	.1518859

Selectivity bias adjustment

In labor market research it is common to include a correction for sample selection bias in the wage equations based on the procedure by Heckman (1976, 1979). Wages are only observed for people who are participating in the labor force and this might be a selective group. The most straightforward approach to account for selection bias in the decomposition is to deduct the selection effects from the overall differential and then apply the standard decomposition formulas to this adjusted differential (Reimers 1983; an alternative approach is followed by Dolton and Makepeace 1986; see Neuman and Oaxaca 2004 for an in-depth treatment of this issue).

If `oaxaca` is used with `heckman`, the decomposition is automatically adjusted for

selection. For example, the following command includes a selection correction in the wage equation for women and decomposes the adjusted wage gap:

```
. oaxaca lnwage educ exper tenure, by(female) model2(heckman, twostep ///
> select(lfp = age agesq married single divorced kids6 kids714))
Blinder-Oaxaca decomposition          Number of obs   =       1434
      1: female = 0
      2: female = 1
```

	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
Differential						
Prediction_1	3.440222	.0174874	196.73	0.000	3.405947	3.474497
Prediction_2	3.275643	.0281554	116.34	0.000	3.220459	3.330827
Difference	.164579	.0331442	4.97	0.000	.0996176	.2295404
Decomposit-n						
Endowments	.0858436	.0157566	5.45	0.000	.0549613	.116726
Coefficients	.0736812	.0311129	2.37	0.018	.0126695	.134693
Interaction	.0050542	.0109895	0.46	0.646	-.0164849	.0265932

Comparing the results to the output in the first example reveals that the uncorrected wages of women are slightly biased downwards (3.267 versus the selectivity corrected 3.276) and the wage gap is somewhat overestimated (0.173 versus the corrected 0.165).

Alternatively, it is sometimes sensible to compute the selection variables outside of `oaxaca` and then use the `adjust()` option to correct the differential. Example:

```
. probit lfp age agesq married single divorced kids6 kids714 if female==1
(output omitted)
. predict xb if e(sample), xb
(759 missing values generated)
. generate mills = normalden(-xb) / (1 - normal(-xb))
(759 missing values generated)
. replace mills = 0 if female==0
(759 real changes made)
. oaxaca lnwage educ exper tenure mills, by(female) adjust(mills)
Blinder-Oaxaca decomposition          Number of obs   =       1434
      1: female = 0
      2: female = 1
```

	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
Differential						
Prediction_1	3.440222	.0174874	196.73	0.000	3.405947	3.474497
Prediction_2	3.266761	.0218659	149.40	0.000	3.223905	3.309618
Difference	.1734607	.0279987	6.20	0.000	.1185843	.2283372
Adjusted	.164579	.033215	4.95	0.000	.0994788	.2296792
Decomposit-n						
Endowments	.0858436	.0157766	5.44	0.000	.0549221	.1167651
Coefficients	.0736812	.0312044	2.36	0.018	.0125217	.1348407
Interaction	.0050542	.0110181	0.46	0.646	-.0165409	.0266493

Using `oaxaca` with nonstandard models

You can also use `oaxaca`, for example, with binary outcome variables and employ a command such as `logit` to estimate the models. You have to understand, however, that `oaxaca` will always apply the decomposition to the linear predictions from the models (based on the first equation if a model contains multiple equations). With `logit` models, for example, the decomposition computed by `oaxaca` is expressed in terms of log odds and not in terms of probabilities or proportions.¹¹

5 References

- Blinder, A. S. 1973. Wage Discrimination: Reduced Form and Structural Estimates. *The Journal of Human Resources* 8: 436–455.
- Cotton, J. 1988. On the Decomposition of Wage Differentials. *The Review of Economics and Statistics* 70: 236–243.
- Daymont, T. N., and P. J. Andrisani. 1984. Job Preferences, College Major, and the Gender Gap in Earnings. *The Journal of Human Resources* 19: 408–428.
- Dolton, P. J., and G. H. Makepeace. 1986. Sample Selection and Male-Female Earnings Differentials in the Graduate Labour Market. *Oxford Economic Papers* 38: 317–341.
- Fairlie, R. W. 2005. An extension of the Blinder-Oaxaca decomposition technique to logit and probit models. *Journal of Economic and Social Measurement* 30: 305–316.
- Fortin, N. M. 2006. Greed, Altruism, and the Gender Wage Gap. University of British Columbia. Available from <http://www.econ.ubc.ca/nfortin/Fortinat8.pdf>.
- Gardeazabal, J., and A. Ugidos. 2004. More on identification in detailed wage decompositions. *The Review of Economics and Statistics* 86: 1034–1036.
- Gomulka, J., and N. Stern. 1990. The Employment of Married Women in the United Kingdom 1970–83. *Economica* 57: 171–199.
- Greene, W. H. 2003. *Econometric Analysis*. 5th ed. Upper Saddle River, NJ: Pearson Education.
- Gujarati, D. N. 2003. *Basic Econometrics*. 4th ed. New York: McGraw-Hill.
- Haisken-DeNew, J. P., and C. M. Schmidt. 1997. Inter-Industry and Inter-Regional Differentials: Mechanics and Interpretation. *The Review of Economics and Statistics* 79: 516–521.

11. Approaches to decompose differences in proportions are provided by, e.g., Gomulka and Stern (1990), Fairlie (2005), or Yun (2005a).

- Hardy, M. A. 1993. *Regression With Dummy Variables*. Newbury Park, CA: Sage.
- Heckman, J. J. 1976. The Common Structure of Statistical Models of Truncation, Sample Selection and Limited Dependent Variables and a Simple Estimator for Such Models. *Annals of Econometrics and Social Measurement* 5: 475–492.
- . 1979. Sample Selection Bias as a Specification Error. *Econometrica* 47: 153–161.
- Heinrichs, J., and P. Kennedy. 2007. A computational trick for calculating the Blinder-Oaxaca decomposition and its standard error. *Economics Bulletin* 3(66): 1–7.
- Hendrickx, J. 1999. dm73: Using categorical variables in Stata. *Stata Technical Bulletin* 52: 2–8.
- Horrace, W. C., and R. L. Oaxaca. 2001. Inter-Industry Wage Differentials and the Gender Wage Gap: An Identification Problem. *Industrial and Labor Relations Review* 54: 611–618.
- Jackson, J. D., and J. T. Lindley. 1989. Measuring the extent of wage discrimination: a statistical test and a caveat. *Applied Economics* 21: 515–540.
- Jann, B. 2003. The Swiss Labor Market Survey 1998 (SLMS 98). *Journal of Applied Social Science Studies* 123: 329–335.
- . 2005a. devcon: Stata module to apply the deviation contrast transform to estimation results. Statistical Software Components S450603. Available from <http://ideas.repec.org/c/boc/bocode/s450603.html>.
- . 2005b. Standard Errors for the Blinder-Oaxaca Decomposition. German Stata Users' Group Meetings 2005. Available from http://repec.org/dsug2005/oaxaca_se_handout.pdf.
- Jones, F. L. 1983. On Decomposing the Wage Gap: A Critical Comment on Blinder's Method. *The Journal of Human Resources* 18: 126–130.
- Jones, F. L., and J. Kelley. 1984. Decomposing Differences Between Groups. A Cautionary Note on Measuring Discrimination. *Sociological Methods and Research* 12: 323–343.
- Kennedy, P. 1986. Interpreting Dummy Variables. *The Review of Economics and Statistics* 68: 174–175.
- Lin, E. S. 2007. On the standard errors of Oaxaca-type decompositions for inter-industry gender wage differentials. *Economics Bulletin* 10(6): 1–11.
- Mood, A. M., F. A. Graybill, and D. C. Boes. 1974. *Introduction to the Theory of Statistics*. New York: McGraw-Hill.
- Neuman, S., and R. L. Oaxaca. 2004. Wage decompositions with selectivity-corrected wage equations: A methodological note. *Journal of Economic Inequality* 2: 3–10.

- Neumark, D. 1988. Employers' Discriminatory Behavior and the Estimation of Wage Discrimination. *The Journal of Human Resources* 23: 279–295.
- Nielsen, H. S. 2000. Wage discrimination in Zambia: an extension of the Oaxaca-Blinder decomposition. *Applied Economics Letters* 7: 405–408.
- Oaxaca, R. 1973. Male-Female Wage Differentials in Urban Labor Markets. *International Economic Review* 14: 693–709.
- Oaxaca, R. L., and M. Ransom. 1998. Calculation of approximate variances for wage decomposition differentials. *Journal of Economic and Social Measurement* 24: 55–61.
- Oaxaca, R. L., and M. R. Ransom. 1994. On discrimination and the decomposition of wage differentials. *Journal of Econometrics* 61: 5–21.
- . 1999. Identification in Detailed Wage Decompositions. *The Review of Economics and Statistics* 81: 154–157.
- O'Donnell, O., E. van Doorslaer, A. Wagstaff, and M. Lindelow. 2008. *Analyzing Health Equity Using Household Survey Data. A Guide to Techniques and Their Implementation*. Washington, DC: The World Bank.
- Polavieja, J. G. 2005. Task Specificity and the Gender Wage Gap. Theoretical Considerations and Empirical Analysis of the Spanish Survey on Wage Structure. *European Sociological Review* 21: 165–181.
- Reimers, C. W. 1983. Labor Market Discrimination Against Hispanic and Black Men. *The Review of Economics and Statistics* 65: 570–579.
- Shrestha, K., and C. Sakellariou. 1996. Wage discrimination: a statistical test. *Applied Economics Letters* 3: 649–651.
- Silber, J., and M. Weber. 1999. Labor market discrimination: are there significant differences between the various decomposition procedures? *Applied Economics* 31: 359–365.
- Stanley, T. D., and S. B. Jarrell. 1998. Gender Wage Discrimination Bias? A Meta-Regression Analysis. *The Journal of Human Resources* 33: 947–973.
- Suits, D. B. 1984. Dummy Variables: Mechanics V. Interpretation. *The Review of Economics and Statistics* 66: 177–180.
- Weesie, J. 1999. sg121: Seemingly unrelated estimation and the cluster-adjusted sandwich estimator. *Stata Technical Bulletin* 52: 34–47.
- Weichselbaumer, D., and R. Winter-Ebmer. 2005. A Meta-Analysis of the International Gender Wage Gap. *Journal of Economic Surveys* 19: 479–511.
- Winsborough, H. H., and P. Dickinson. 1971. Components of Negro-White Income Differences. In *Proceedings of the Social Statistics Section*, 6–8. Washington, DC: American Statistical Association.

Yun, M.-S. 2005a. Hypothesis tests when decomposing differences in the first moment. *Journal of Economic and Social Measurement* 30: 295–304.

———. 2005b. A Simple Solution to the Identification Problem in Detailed Wage Decompositions. *Economic Inquiry* 43: 766–772.